




Met Polyclonal Antibody

| | |
|---------------------------------|--|
| Swiss-Prot No.: | P08581 |
| Molecular Weight: | 155.527 |
| Applications: | WB,IF,ELISA |
| Reactivity: | Human,Mouse,Rat |
| Purification: | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Immunogen: | Synthesized peptide derived from human Met around the non-phosphorylation site of Y1349. |
| Specificity/Sensitivity: | Met Polyclonal Antibody detects endogenous levels of Met protein. |
| Other Names: | MET; Hepatocyte growth factor receptor; HGF receptor; HGF/SF receptor; Proto-oncogene c-Met; Scatter factor receptor; SF receptor; Tyrosine-protein kinase Met |
| Storage/Stability: | -20°C/1 year |
| Source: | Rabbit |
| Form of Antibody: | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Protein Concentration: | 1 mg/ml |

For Research Use Only

| | |
|-------------------|---|
| |  |
| | Western Blot analysis of various cells using Met Polyclonal Antibody |
| Gene Name: | MET |
| Dilution: | Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications. |