




NF-YC Polyclonal Antibody

| | |
|---------------------------------|---|
| Swiss-Prot No.: | Q13952 |
| Molecular Weight: | 50.302 |
| Applications: | WB,IHC,ELISA |
| Reactivity: | Human,Mouse,Rat |
| Purification: | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Immunogen: | Synthesized peptide derived from the N-terminal region of human NF-YC. at AA range: 10-90 |
| Specificity/Sensitivity: | NF-YC Polyclonal Antibody detects endogenous levels of NF-YC protein. |
| Other Names: | NFYC; Nuclear transcription factor Y subunit gamma; CAAT box DNA-binding protein subunit C; Nuclear transcription factor Y subunit C; NF-YC; Transactivator HSM-1/2 |
| Storage/Stability: | -20°C/1 year |
| Source: | Rabbit |
| Form of Antibody: | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Protein Concentration: | 1 mg/ml |

For Research Use Only

| | |
|-------------------|---|
| |  <p>(kD)</p> <p>117-</p> <p>85-</p> <p>48- NF-YC</p> <p>34-</p> <p>26-</p> <p>19-</p> <p>3T3</p> |
| | Western Blot analysis of various cells using NF-YC Polyclonal Antibody |
| Gene Name: | NFYC |
| Dilution: | Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications. |