




## NIPA Polyclonal Antibody

<b>Swiss-Prot No.:</b>	Q86WB0
<b>Molecular Weight:</b>	55.262
<b>Applications:</b>	WB,IF,ELISA
<b>Reactivity:</b>	Human,Mouse,Rat,Monkey
<b>Purification:</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Immunogen:</b>	Synthesized peptide derived from human NIPA around the non-phosphorylation site of S354.
<b>Specificity/Sensitivity:</b>	NIPA Polyclonal Antibody detects endogenous levels of NIPA protein.
<b>Other Names:</b>	ZC3HC1; NIPA; HSPC216; Nuclear-interacting partner of ALK; Nuclear-interacting partner of anaplastic lymphoma kinase; hNIPA; Zinc finger C3HC-type protein 1
<b>Storage/Stability:</b>	-20°C/1 year
<b>Source:</b>	Rabbit
<b>Form of Antibody:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Protein Concentration:</b>	1 mg/ml

For Research Use Only

	 <p>Western blot analysis of COS7 cells. The blot shows a single band at approximately 55 kD. Molecular weight markers are indicated on the left: 117, 85, 48, 34, 26, and 19 kD.</p>
	Western Blot analysis of various cells using NIPA Polyclonal Antibody
Gene Name:	ZC3HC1
Dilution:	Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.