




## Op18 Polyclonal Antibody

<b>Swiss-Prot No.:</b>	P16949
<b>Molecular Weight:</b>	17.303
<b>Applications:</b>	WB,IHC,ELISA
<b>Reactivity:</b>	Human,Mouse,Rat
<b>Purification:</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Immunogen:</b>	Synthesized peptide derived from human Op18 around the non-phosphorylation site of S38.
<b>Specificity/Sensitivity:</b>	Op18 Polyclonal Antibody detects endogenous levels of Op18 protein.
<b>Other Names:</b>	STMN1; C1orf215; LAP18; OP18; Stathmin; Leukemia-associated phosphoprotein p18; Metablastin; Oncoprotein 18; Op18; Phosphoprotein p19; pp19; Prosolin; Protein Pr22; pp17
<b>Storage/Stability:</b>	-20°C/1 year
<b>Source:</b>	Rabbit
<b>Form of Antibody:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Protein Concentration:</b>	1 mg/ml

For Research Use Only

	 <p>(kD)</p> <p>117- 85- 48- 34- 26- 19-</p> <p>Jurkat</p> <p>Op18</p>
	Western Blot analysis of various cells using Op18 Polyclonal Antibody
Gene Name:	STMN1
Dilution:	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications.