




## p18 Polyclonal Antibody

<b>Swiss-Prot No.:</b>	P42773
<b>Molecular Weight:</b>	18.127
<b>Applications:</b>	WB,IHC,ELISA
<b>Reactivity:</b>	Human,Mouse,Monkey
<b>Purification:</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Immunogen:</b>	Synthesized peptide derived from the C-terminal region of human p18. at AA rangle: 80-160
<b>Specificity/Sensitivity:</b>	p18 Polyclonal Antibody detects endogenous levels of p18 protein.
<b>Other Names:</b>	CDKN2C; CDKN6; Cyclin-dependent kinase 4 inhibitor C; Cyclin-dependent kinase 6 inhibitor; p18-INK4c; p18-INK6
<b>Storage/Stability:</b>	-20°C/1 year
<b>Source:</b>	Rabbit
<b>Form of Antibody:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Protein Concentration:</b>	1 mg/ml

	 <p>Western blot analysis of COS7 cells using p18 Polyclonal Antibody. The blot shows a single band at approximately 19 kDa. Molecular weight markers (kDa) are indicated on the left: 117, 85, 48, 34, 26, 19.</p>
	Western Blot analysis of various cells using p18 Polyclonal Antibody
<b>Gene Name:</b>	CDKN2C
<b>Dilution:</b>	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications.