



PSK2 Polyclonal Antibody

| | |
|---------------------------------|--|
| Swiss-Prot No.: | Q7L7X3 |
| Molecular Weight: | 116.07 |
| Applications: | WB,IHC,ELISA |
| Reactivity: | Human,Mouse,Rat |
| Purification: | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Immunogen: | Synthesized peptide derived from the Internal region of human PSK2. at AA range: 400-480 |
| Specificity/Sensitivity: | PSK2 Polyclonal Antibody detects endogenous levels of PSK2 protein. |
| Other Names: | TAOK1; KIAA1361; MAP3K16; MARKK; Serine/threonine-protein kinase TAO1; Kinase from chicken homolog B; hKFC-B; MARK Kinase; MARKK; Prostate-derived sterile 20-like kinase 2; PSK-2; PSK2; Prostate-derived STE20-like kinase 2; Thousand and one |
| Storage/Stability: | -20°C/1 year |
| Source: | Rabbit |
| Form of Antibody: | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |

For Research Use Only

| | |
|-------------------------------|---|
| Protein Concentration: | 1 mg/ml |
| Gene Name: | TAOK1 |
| Dilution: | Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications. |