




## PSMD2 Polyclonal Antibody

<b>Swiss-Prot No.:</b>	Q13200
<b>Molecular Weight:</b>	100.2
<b>Applications:</b>	WB,IHC,ELISA
<b>Reactivity:</b>	Human,Mouse,Rat
<b>Purification:</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Immunogen:</b>	Synthesized peptide derived from the N-terminal region of human PSMD2. at AA range: 70-150
<b>Specificity/Sensitivity:</b>	PSMD2 Polyclonal Antibody detects endogenous levels of PSMD2 protein.
<b>Other Names:</b>	PSMD2; TRAP2; 26S proteasome non-ATPase regulatory subunit 2; 26S proteasome regulatory subunit RPN1; 26S proteasome regulatory subunit S2; 26S proteasome subunit p97; Protein 55.11; Tumor necrosis factor type 1 receptor-associated protein
<b>Storage/Stability:</b>	-20°C/1 year
<b>Source:</b>	Rabbit
<b>Form of Antibody:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

**For Research Use Only**

<b>Protein Concentration:</b>	1 mg/ml
	<p>(kD) Jurkat</p>  <p>117- PSMD2 85- 48- 34- 26- 19-</p>
	Western Blot analysis of various cells using PSMD2 Polyclonal Antibody
<b>Gene Name:</b>	PSMD2
<b>Dilution:</b>	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications.