




## SDHA Polyclonal Antibody

<b>Swiss-Prot No.:</b>	P31040
<b>Molecular Weight:</b>	72.692
<b>Applications:</b>	WB,ELISA
<b>Reactivity:</b>	Human,Mouse,Rat
<b>Purification:</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Immunogen:</b>	Synthesized peptide derived from the C-terminal region of human SDHA.
<b>Specificity/Sensitivity:</b>	SDHA Polyclonal Antibody detects endogenous levels of SDHA protein.
<b>Other Names:</b>	SDHA; SDH2; SDHF; Succinate dehydrogenase [ubiquinone] flavoprotein subunit; mitochondrial; Flavoprotein subunit of complex II; Fp
<b>Storage/Stability:</b>	-20°C/1 year
<b>Source:</b>	Rabbit
<b>Form of Antibody:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Protein Concentration:</b>	1 mg/ml

	 <p>Western blot analysis of various cells using SDHA Polyclonal Antibody. The blot shows bands for SDHA in 3T3, Jurkat, K562, HepG2, and Rat kidney samples. Molecular weight markers are indicated on the left: 120, 90, 50, 39, and 27 kDa. The SDHA band is located between 90 and 120 kDa.</p>
	Western Blot analysis of various cells using SDHA Polyclonal Antibody
<b>Gene Name:</b>	SDHA
<b>Dilution:</b>	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.