



|  |                                       |  |
|--|---------------------------------------|--|
|  | <h1>Skp1 p19 Polyclonal Antibody</h1> |  |
|--|---------------------------------------|--|

|                                 |  |
|---------------------------------|--|
| <b>Swiss-Prot No.:</b>          | P63208   |
| <b>Molecular Weight:</b>        | 18.658   |
| <b>Applications:</b>            | WB,IHC,ELISA   |
| <b>Reactivity:</b>              | Human,Mouse,Rat,Monkey   |
| <b>Purification:</b>            | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  |
| <b>Immunogen:</b>               | Synthesized peptide derived from the Internal region of human Skp1 p19. at AA range: 10-90   |
| <b>Specificity/Sensitivity:</b> | Skp1 p19 Polyclonal Antibody detects endogenous levels of Skp1 p19 protein.  |
| <b>Other Names:</b>             | SKP1; EMC19; OCP2; SKP1A; TCEB1L; S-phase kinase-associated protein 1; Cyclin-A/CDK2-associated protein p19; Organ of Corti protein 2; OCP-2; Organ of Corti protein II; OCP-II; RNA polymerase II elongation factor-like protein; SIII; Transcr |
| <b>Storage/Stability:</b>       | -20°C/1 year   |
| <b>Source:</b>                  | Rabbit   |
| <b>Form of Antibody:</b>        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| <b>Protein Concentration:</b>   | 1 mg/ml  |

**For Research Use Only**

|            |   |
|------------|---|
|            |  <p>Western blot analysis of COS7 cells using Skp1 p19 Polyclonal Antibody. The blot shows a single band at approximately 19 kD.</p> |
|            | Western Blot analysis of various cells using Skp1 p19 Polyclonal Antibody   |
| Gene Name: | SKP1  |
| Dilution:  | Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications.   |