
	<h1>Smad2 Polyclonal Antibody</h1>	
--	------------------------------------	--

<b>Swiss-Prot No.:</b>	Q15796
<b>Molecular Weight:</b>	52.306
<b>Applications:</b>	WB,IHC,IF,ELISA
<b>Reactivity:</b>	Human,Mouse,Rat
<b>Purification:</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Immunogen:</b>	Synthesized peptide derived from human Smad2 around the non-phosphorylation site of S467.
<b>Specificity/Sensitivity:</b>	Smad2 Polyclonal Antibody detects endogenous levels of Smad2 protein.
<b>Other Names:</b>	SMAD2; MADH2; MADR2; Mothers against decapentaplegic homolog 2; MAD homolog 2; Mothers against DPP homolog 2; JV18-1; Mad-related protein 2; hMAD-2; SMAD family member 2; SMAD 2; Smad2; hSMAD2
<b>Storage/Stability:</b>	-20°C/1 year
<b>Source:</b>	Rabbit
<b>Form of Antibody:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Protein Concentration:</b>	1 mg/ml

**For Research Use Only**

	 <p>COLO</p> <p>170— 130— 100— 70— 55— 40— 35— 25— 15—</p> <p>Smad2 (Ab-467)</p>
	Western Blot analysis of COLO cells using Smad2 Polyclonal Antibody
<b>Gene Name:</b>	SMAD2
<b>Dilution:</b>	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.