




## STAG3 Polyclonal Antibody

<b>Swiss-Prot No.:</b>	Q9UJ98
<b>Molecular Weight:</b>	139.034
<b>Applications:</b>	WB,IHC,ELISA
<b>Reactivity:</b>	Human
<b>Purification:</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Immunogen:</b>	Synthesized peptide derived from the C-terminal region of human STAG3. at AA range: 1140-1220
<b>Specificity/Sensitivity:</b>	STAG3 Polyclonal Antibody detects endogenous levels of STAG3 protein.
<b>Other Names:</b>	STAG3; Cohesin subunit SA-3; SCC3 homolog 3; Stromal antigen 3; Stromalin-3
<b>Storage/Stability:</b>	-20°C/1 year
<b>Source:</b>	Rabbit
<b>Form of Antibody:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Protein Concentration:</b>	1 mg/ml

	 <p>Western blot analysis of Jurkat cells using STAG3 Polyclonal Antibody. The blot shows a single band at approximately 130 kDa, labeled STAG3. Molecular weight markers are indicated on the left: 170, 130, 100, 70, 55, 40, 35, 25, and 15 kDa.</p>
	Western Blot analysis of Jurkat cells using STAG3 Polyclonal Antibody
<b>Gene Name:</b>	STAG3
<b>Dilution:</b>	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications.