



STEAP2 Polyclonal Antibody

| | |
|---------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Swiss-Prot No.: | Q8NFT2 |
| Molecular Weight: | 55.994 |
| Applications: | IHC, IF, ELISA |
| Reactivity: | Human, Mouse |
| Purification: | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Immunogen: | Synthesized peptide derived from the C-terminal region of human STEAP2. at AA range: 400-480 |
| Specificity/Sensitivity: | STEAP2 Polyclonal Antibody detects endogenous levels of STEAP2 protein. |
| Other Names: | STEAP2; PCANAP1; STAMP1; Metalloreductase STEAP2; Prostate cancer-associated protein 1; Protein up-regulated in metastatic prostate cancer; PUMPCn; Six-transmembrane epithelial antigen of prostate 2; SixTransMembrane protein of prostate 1 |
| Storage/Stability: | -20°C/1 year |
| Source: | Rabbit |
| Form of Antibody: | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |

| | |
|-------------------------------|--------------------------------------------------------------------------------------------------------------------------------|
| Protein Concentration: | 1 mg/ml |
| Gene Name: | STEAP2 |
| Dilution: | Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications. |