



## TACC1 Polyclonal Antibody

|                                 |   |
|---------------------------------|---|
| <b>Swiss-Prot No.:</b>          | 075410  |
| <b>Molecular Weight:</b>        | 87.794  |
| <b>Applications:</b>            | WB,IHC,IF,ELISA   |
| <b>Reactivity:</b>              | Human,Mouse   |
| <b>Purification:</b>            | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| <b>Immunogen:</b>               | Synthesized peptide derived from the N-terminal region of human TACC1. at AA range: 30-110                            |
| <b>Specificity/Sensitivity:</b> | TACC1 Polyclonal Antibody detects endogenous levels of TACC1 protein.   |
| <b>Other Names:</b>             | TACC1; KIAA1103; Transforming acidic coiled-coil-containing protein 1; Gastric cancer antigen Ga55; Taxin-1           |
| <b>Storage/Stability:</b>       | -20°C/1 year  |
| <b>Source:</b>                  | Rabbit  |
| <b>Form of Antibody:</b>        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| <b>Protein Concentration:</b>   | 1 mg/ml   |
| <b>Gene Name:</b>               | TACC1   |

For Research Use Only

|                  |  |
|------------------|--|
| <b>Dilution:</b> | Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications. |
|------------------|--|