




TBC1D3A/B/C Polyclonal Antibody

| | |
|---------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Swiss-Prot No.: | Q8IZP1/A6NDS4/Q6IPX1 |
| Molecular Weight: | 62.199 |
| Applications: | WB,IF,ELISA |
| Reactivity: | Human |
| Purification: | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Immunogen: | Synthesized peptide derived from the C-terminal region of human TBC1D3A/B/C. at AA range: 450-530 |
| Specificity/Sensitivity: | TBC1D3A/B/C Polyclonal Antibody detects endogenous levels of TBC1D3A/B/C protein. |
| Other Names: | TBC1D3; PRC17; TBC1D3A; TBC1 domain family member 3; Prostate cancer gene 17 protein; Protein TRE17-alpha; Rab GTPase-activating protein PRC17; TBC1D3B; TBC1D3I; TBC1 domain family member 3B; TBC1D3C; TBC1D3D; TBC1 domain family member 3C |
| Storage/Stability: | -20°C/1 year |
| Source: | Rabbit |
| Form of Antibody: | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |

For Research Use Only

| | |
|-------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Protein Concentration: | 1 mg/ml |
| | <p style="text-align: center;">HeLa</p>  <p>(kD)</p> <p>117-</p> <p>85-</p> <p>48-</p> <p>34-</p> <p>26-</p> <p>19-</p> |
| | Western Blot analysis of various cells using TBC1D3A/B/C Polyclonal Antibody |
| Gene Name: | TBC1D3 |
| Dilution: | Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications. |