




## TdT Polyclonal Antibody

<b>Swiss-Prot No.:</b>	P04053
<b>Molecular Weight:</b>	58.437
<b>Applications:</b>	WB,ELISA
<b>Reactivity:</b>	Human
<b>Purification:</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Immunogen:</b>	Synthesized peptide derived from the Internal region of human TdT. at AA range: 350-430
<b>Specificity/Sensitivity:</b>	TdT Polyclonal Antibody detects endogenous levels of TdT protein.
<b>Other Names:</b>	DNTT; TDT; DNA nucleotidylexotransferase; Terminal addition enzyme; Terminal deoxynucleotidyltransferase; Terminal transferase
<b>Storage/Stability:</b>	-20°C/1 year
<b>Source:</b>	Rabbit
<b>Form of Antibody:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Protein Concentration:</b>	1 mg/ml

For Research Use Only

	 <p>HepG2</p> <p>170— 130— 100— 70— 55— TdT 40— 35— 25— 15—</p>
	Western Blot analysis of HepG2 cells using TdT Polyclonal Antibody
<b>Gene Name:</b>	DNTT
<b>Dilution:</b>	Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.