


<b>Swiss-Prot No.:</b>	Q8WUA4
<b>Molecular Weight:</b>	100.68
<b>Applications:</b>	WB,IHC,ELISA
<b>Reactivity:</b>	Human
<b>Purification:</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Immunogen:</b>	Synthesized peptide derived from the N-terminal region of human TFIIIC110. at AA range: 40-120
<b>Specificity/Sensitivity:</b>	TFIIIC110 Polyclonal Antibody detects endogenous levels of TFIIIC110 protein.
<b>Other Names:</b>	GTF3C2; KIAA0011; General transcription factor 3C polypeptide 2; TF3C-beta; Transcription factor IIIC 110 kDa subunit; TFIIIC 110 kDa subunit; TFIIIC110; Transcription factor IIIC subunit beta
<b>Storage/Stability:</b>	-20°C/1 year
<b>Source:</b>	Rabbit
<b>Form of Antibody:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Protein Concentration:</b>	1 mg/ml

**For Research Use Only**

	 <p>Western blot analysis of HepG2 cells using TFIIC110 Polyclonal Antibody. The blot shows a single band at approximately 110 kD. Molecular weight markers are indicated on the left at 170, 130, 95, 72, and 55 kD.</p>
	Western Blot analysis of various cells using TFIIC110 Polyclonal Antibody
Gene Name:	GTF3C2
Dilution:	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications.