

TNF-α Polyclonal Antibody

| 0.1.0.11 | D0 4 0 7 7 |
|--------------------------|--|
| Swiss-Prot No.: | P01375 |
| Molecular Weight: | 25.644 |
| Applications: | WB,IHC,IF,ELISA |
| Reactivity: | Human, Mouse, Rat |
| Purification: | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Immunogen: | Synthesized peptide derived from the Internal region of human TNF- α . at AA rangle: 110-190 |
| Specificity/Sensitivity: | TNF- α Polyclonal Antibody detects endogenous levels of TNF- α protein. |
| Other Names: | TNF; TNFA; TNFSF2; Tumor necrosis factor; Cachectin; TNF- alpha; Tumor necrosis factor ligand superfamily member 2; TNF- a |
| Storage/Stability: | -20°C/1 year |
| Source: | Rabbit |
| Form of Antibody: | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Protein Concentration: | 1 mg/ml |

| | (kD) Jurkat 117- 85- 48- 34- 26- 19- TNF-α |
|------------|--|
| | Western Blot analysis of various cells using TNF-α Polyclonal Antibody |
| Gene Name: | TNF |
| Dilution: | Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications. |