




## TSLC1 Polyclonal Antibody

|                                 |  |
|---------------------------------|--|
| <b>Swiss-Prot No.:</b>          | Q9BY67   |
| <b>Molecular Weight:</b>        | 48.509   |
| <b>Applications:</b>            | WB,IF,ELISA  |
| <b>Reactivity:</b>              | Human,Mouse  |
| <b>Purification:</b>            | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  |
| <b>Immunogen:</b>               | Synthesized peptide derived from the C-terminal region of human TSLC1. at AA range: 370-450  |
| <b>Specificity/Sensitivity:</b> | TSLC1 Polyclonal Antibody detects endogenous levels of TSLC1 protein.  |
| <b>Other Names:</b>             | CADM1; IGSF4; IGSF4A; NECL2; SYNCAM; TSLC1; Cell adhesion molecule 1; Immunoglobulin superfamily member 4; IgSF4; Nectin-like protein 2; NECL-2; Spermatogenic immunoglobulin superfamily; SgIgSF; Synaptic cell adhesion molecule; SynCAM; Tumo |
| <b>Storage/Stability:</b>       | -20°C/1 year   |
| <b>Source:</b>                  | Rabbit   |
| <b>Form of Antibody:</b>        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |

For Research Use Only

|                               |   |
|-------------------------------|---|
| <b>Protein Concentration:</b> | 1 mg/ml   |
|                               | <p style="text-align: center;">HeLa</p>  <p>(kD)</p> <p>117-</p> <p>85-</p> <p>48-</p> <p>34-</p> <p>26-</p> <p>19-</p> |
|                               | Western Blot analysis of various cells using TSLC1 Polyclonal Antibody  |
| <b>Gene Name:</b>             | CADM1   |
| <b>Dilution:</b>              | Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.   |