

	<h2>VPAC2 Polyclonal Antibody</h2>	
--	------------------------------------	--

<b>Swiss-Prot No.:</b>	P41587
<b>Molecular Weight:</b>	49.479
<b>Applications:</b>	WB,IHC,IF,ELISA
<b>Reactivity:</b>	Human,Mouse,Rat
<b>Purification:</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Immunogen:</b>	Synthesized peptide derived from the Internal region of human VPAC2. at AA range: 50-130
<b>Specificity/Sensitivity:</b>	VPAC2 Polyclonal Antibody detects endogenous levels of VPAC2 protein.
<b>Other Names:</b>	VIPR2; VIP2R; Vasoactive intestinal polypeptide receptor 2; VIP-R-2; Helodermin-preferring VIP receptor; Pituitary adenylate cyclase-activating polypeptide type III receptor; PACAP type III receptor; PACAP-R-3; PACAP-R3; VPAC2
<b>Storage/Stability:</b>	-20°C/1 year
<b>Source:</b>	Rabbit
<b>Form of Antibody:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Protein Concentration:</b>	1 mg/ml

**For Research Use Only**

<b>Gene Name:</b>	VIPR2
<b>Dilution:</b>	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.