




## XRCC2 Polyclonal Antibody

|                                 |   |
|---------------------------------|---|
| <b>Swiss-Prot No.:</b>          | 043543  |
| <b>Molecular Weight:</b>        | 31.956  |
| <b>Applications:</b>            | WB,IHC,ELISA  |
| <b>Reactivity:</b>              | Human   |
| <b>Purification:</b>            | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| <b>Immunogen:</b>               | Synthesized peptide derived from the C-terminal region of human XRCC2. at AA rangle: 180-260                          |
| <b>Specificity/Sensitivity:</b> | XRCC2 Polyclonal Antibody detects endogenous levels of XRCC2 protein.   |
| <b>Other Names:</b>             | XRCC2; DNA repair protein XRCC2; X-ray repair cross-complementing protein 2   |
| <b>Storage/Stability:</b>       | -20°C/1 year  |
| <b>Source:</b>                  | Rabbit  |
| <b>Form of Antibody:</b>        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| <b>Protein Concentration:</b>   | 1 mg/ml   |

|            |  |
|------------|--|
|            |   |
|            | Western Blot analysis of various cells using XRCC2 Polyclonal Antibody   |
| Gene Name: | XRCC2  |
| Dilution:  | Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications. |