




## IGF-IIR Polyclonal Antibody

<b>Swiss-Prot No.:</b>	P11717
<b>Molecular Weight:</b>	274.276
<b>Applications:</b>	WB,ELISA
<b>Reactivity:</b>	Human
<b>Purification:</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Immunogen:</b>	Synthesized peptide derived from the C-terminal region of human IGF-IIR.
<b>Specificity/Sensitivity:</b>	IGF-IIR Polyclonal Antibody detects endogenous levels of IGF-IIR protein.
<b>Other Names:</b>	IGF2R; MPRI; Cation-independent mannose-6-phosphate receptor; CI Man-6-P receptor; CI-MPR; M6PR; 300 kDa mannose 6-phosphate receptor; MPR 300;Insulin-like growth factor 2 receptor; Insulin-like growth factor II receptor; IGF-II receptor; M6P/IGF2 recepto
<b>Storage/Stability:</b>	-20°C/1 year
<b>Source:</b>	Rabbit
<b>Form of Antibody:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

**For Research Use Only**

<b>Protein Concentration:</b>	1 mg/ml
	 <p>Western blot analysis of MCF7 cells. The image shows a single prominent band at approximately 270 kDa, labeled 'MCF7'. The molecular weight markers on the left are 300, 250, 180, 130, 100, 70, and 50 kDa.</p>
	Western Blot analysis of MCF7 cells using IGF-IIR Polyclonal Antibody.
<b>Gene Name:</b>	IGF2R
<b>Dilution:</b>	Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.