


	<h2 style="margin: 0;">RANKL Polyclonal Antibody</h2>
--	---

<b>Swiss-Prot No.:</b>	014788
<b>Molecular Weight:</b>	35.478
<b>Applications:</b>	WB,IHC,ELISA
<b>Reactivity:</b>	Human,Mouse,Rat
<b>Purification:</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Immunogen:</b>	Synthesized peptide derived from the C-terminal region of human RANKL.
<b>Specificity/Sensitivity:</b>	RANKL Polyclonal Antibody detects endogenous levels of RANKL protein.
<b>Other Names:</b>	TNFSF11; OPGL; RANKL; TRANCE; Tumor necrosis factor ligand superfamily member 11; Osteoclast differentiation factor; ODF; Osteoprotegerin ligand; OPGLReceptor activator of nuclear factor kappa-B ligand; RANKL; TNF-related activation-induced cytokine; TRAN
<b>Storage/Stability:</b>	-20°C/1 year
<b>Source:</b>	Rabbit
<b>Form of Antibody:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

**For Research Use Only**

<b>Protein Concentration:</b>	1 mg/ml
	
	Western Blot analysis of 293 cells using RANKL Polyclonal Antibody.
<b>Gene Name:</b>	TNFSF11
<b>Dilution:</b>	Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.