




NCoA-3 Polyclonal Antibody

| | |
|---------------------------------|---|
| Swiss-Prot No.: | Q9Y6Q9 |
| Molecular Weight: | 155.293 |
| Applications: | WB,IHC,ELISA |
| Reactivity: | Human, Mouse, Rat |
| Purification: | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Immunogen: | Synthesized peptide derived from the Internal region of human NCoA-3. |
| Specificity/Sensitivity: | NCoA-3 Polyclonal Antibody detects endogenous levels of NCoA-3 protein. |
| Other Names: | NCOA3; AIB1; BHLHE42; RAC3; TRAM1; Nuclear receptor coactivator 3; NCoA-3; ACTR; Amplified in breast cancer 1 protein; AIB-1; CBP-interacting protein; pCIP; Class E basic helix-loop-helix protein 42; bHLHe42; Receptor-associated coactivator 3; RAC-3; Ster |
| Storage/Stability: | -20°C/1 year |
| Source: | Rabbit |
| Form of Antibody: | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |

For Research Use Only

| | |
|-------------------------------|---|
| Protein Concentration: | 1 mg/ml |
| |  |
| | Western Blot analysis of HeLa, HepG2 cells using NCoA-3 Polyclonal Antibody. |
| Gene Name: | NCOA3 |
| Dilution: | Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications. |