




## VPAC1 Polyclonal Antibody

<b>Swiss-Prot No.:</b>	P32241
<b>Molecular Weight:</b>	51.547
<b>Applications:</b>	WB ELISA
<b>Reactivity:</b>	Human
<b>Purification:</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Immunogen:</b>	Synthesized peptide derived from the Internal region of human VPAC1.
<b>Specificity/Sensitivity:</b>	VPAC1 Polyclonal Antibody detects endogenous levels of VPAC1 protein.
<b>Other Names:</b>	VIPR1; Vasoactive intestinal polypeptide receptor 1; VIP-R-1; Pituitary adenylate cyclase-activating polypeptide type II receptor; PACAP type II receptor; PACAP-R-2; PACAP-R2; VPAC1
<b>Storage/Stability:</b>	-20°C/1 year
<b>Source:</b>	Rabbit
<b>Form of Antibody:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Protein Concentration:</b>	1 mg/ml

For Research Use Only

	
	Western Blot analysis of HepG2 cells using VPAC1 Polyclonal Antibody.
<b>Gene Name:</b>	VIPR1
<b>Dilution:</b>	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.