



## ABL1 Polyclonal Antibody

E90282

**Catalog Number:** E90282**Amount:** 100ul

**Background:** The c-Abl proto-oncogene encodes a nonreceptor protein tyrosine kinase that is ubiquitously expressed and highly conserved in metazoan evolution. c-Abl protein is distributed in both the nucleus and the cytoplasm of cells. It is implicated in regulating cell proliferation, differentiation, apoptosis, cell adhesion and stress responses (1-3). c-Abl kinase activity is increased in vivo by diverse physiological stimuli including integrin activation, PDGF stimulation and binding to c-Jun, Nck and RFX1 (2,4). The in vivo mechanism of regulation of c-Abl kinase activity is not completely understood. Tyr245 is located in the linker region between the SH2 and catalytic domains. This positioning is conserved among Abl family members. Phosphorylation of Tyr245 is involved in the activation of c-Abl kinase (5). In addition, phosphorylation of Tyr412, which is located in the kinase activation loop of c-Abl, is required for kinase activity (6).

**Species:** Rabbit**Isotype:** IgG

**Storage/Stability:** Store at -20oC or -80oC. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonyms:** ABL1;ABL;JTK7;bcr/abl;c-ABL;p150;v-abl ;

**Immunogen:** C term -peptide of human ABL1

**Purification:** Affinity purification

**Reactivity:** H M R

**Applications:** WB

**Molecular Weight:** 123kDa

**Swiss-Prot No.:** P00519

**Gene ID:** 25

**References:** 1. Wang, J.Y. et al. (2000) Oncogene 19, 5643-5650. 2. Van Etten, R.A. et al. (1999) Trends Cell. Biol. 9, 179-182. 3. Danial, N.N. et al. (2000) Oncogene 19, 2523-2531. 4. Shaul, Y. et al. (2000) Cell Death Differ. 7, 10-16. 5. Brasher, B.B. et al. (2000) J. Biol. Chem. 275, 35631-35637. 6. Pluk, H. et al. (2002) Cell 108, 247-259.

**For Research Use Only**

