

MALT1Polyclonal Antibody

| Amount: 100ul Restaround: Museus essesiated lumphoid tissue lumphome translocation protein 1 belongs | | | | |
|--|---|--|--|--|
| Packground: Musees associated hyperbold tissue hyperbolic translocation protein 1 belongs | | | | |
| Background: Mucosa-associated lymphoid tissue lymphoma translocation protein 1 belongs | 0515 | | | |
| caspase-like protein superfamily.MALT1 plays a potential role in the regulation of apopt and induces NFKB activation and required for the normal development of B cells and for | · tha | | | |
| | activation of the IKK complex. It acts as a caspase and mediates additional functions related | | | |
| | to T cell activation, cleaving both the ubiquitin editing enzyme,TNFAIP3,and the C terminus | | | |
| | of BCL10. It has two isoforms. Researches showed that inhibition of MALT1 blocks A20 and | | | |
| BCL10 cleavage, reduces NFKB activity, and decreases the expression of NFKB tar | BCL10 cleavage, reduces NFKB activity, and decreases the expression of NFKB targets | | | |
| BCLXL. | 0 | | | |
| Species: Rabbit | | | | |
| Isotype: IgG | | | | |
| Storage/Stability: Store at -20oC or -80oC. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium az | Store at -20oC or -80oC. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, | | | |
| 50% glycerol, pH7.3. | | | | |
| Synonyms: DKFZp434L132; MLT; MLT1 | DKFZp434L132; MLT; MLT1 | | | |
| Immunogen: Recombinant protein of humanMALT1 | Recombinant protein of humanMALT1 | | | |
| Purification: Affinity purification | | | | |
| Reactivity: HM R | | | | |
| | | | | |
| | 92kDa | | | |
| | Q9UDY8 | | | |
| Gene ID: 10892 | | | | |

| IHC 1:50 - 1:200 | | | |
|------------------|----------------|--|--|
| | IF 1:20 - 1:50 | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |

For Research Use Only