

HGF Polyclonal Antibody

Catalog Number: E91193

Amount: 100ul

Background: Hepatocyte growth factor regulates cell growth, cell motility, and morphogenesis by

activating a tyrosine kinase signaling cascade after binding to the proto-oncogenic c-Met receptor. Hepatocyte growth factor is secreted by mesenchymal cells and acts as a multi-functional cytokine on cells of mainly epithelial origin. Its ability to stimulate mitogenesis, cell motility, and matrix invasion gives it a central role in angiogenesis, tumorogenesis, and tissue regeneration. It is secreted as a single inactive polypeptide and is cleaved by serine proteases into a 69-kDa alpha-chain and 34-kDa beta-chain. A disulfide bond between the alpha and beta chains produces the active, heterodimeric molecule. The protein belongs to the plasminogen subfamily of S1 peptidases but has no detectable protease activity. Alternative splicing of this gene produces multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008]

Species: Rabbit Isotype: IgG


Storage/Stability: Store at -20oC or -80oC. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,

50% glycerol, pH7.3.

Synonyms: SF; HGFB; HPTA; F-TCF; DFNB39 **Immunogen:** Recombinant proteinof human HGF

Purification: Affinity purification

Reactivity: H M R
Applications: WB IHC
Molecular Weight: 83kDa
Swiss-Prot No.: P14210
Gene ID: 3082



For Research Use Only

Support: service@enogene.com

Order: order@enogene.com

Web: www.enogene.com