

KIR3DL1 Polyclonal Antibody

Catalog Number: E91617 Amount: 100ul

> Killer cell immunoglobulin-like receptors (KIRs) are transmembrane glycoproteins Background:

expressed by natural killer cells and subsets of T cells. The KIR genes are polymorphic and highly homologous and they are found in a cluster on chromosome 19q13.4 within the 1 Mb leukocyte receptor complex (LRC). The gene content of the KIR gene cluster varies among haplotypes, although several "framework" genes are found in all haplotypes (KIR3DL3, KIR3DP1, KIR3DL4, KIR3DL2). The KIR proteins are classified by the number of extracellular immunoglobulin domains (2D or 3D) and by whether they have a long (L) or short (S) cytoplasmic domain. KIR proteins with the long cytoplasmic domain transduce inhibitory signals upon ligand binding via an immune tyrosine-based inhibitory motif (ITIM), while KIR proteins with the short cytoplasmic domain lack the ITIM motif and instead associate with the TYRO protein tyrosine kinase binding protein to transduce activating signals. The ligands for several KIR proteins are subsets of HLA class I molecules; thus, KIR proteins are thought to play an important role in regulation of the immune response.

Rabbit Species: Isotype: IgG

Storage/Stability: Store at -20oC or -80oC. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,

50% glycerol, pH7.3.


Synonyms: KIR; NKB1; NKAT3; NKB1B; CD158E1; KIR3DL1/S1;

Immunogen: Recombinant protein of human KIR3DL1

Purification: Affinity purification

Reactivity:

Applications: WB IHC Molecular Weight: 49kDa Swiss-Prot No.: P43629 Gene ID:



For Research Use Only

Support: service@enogene.com

Order: order@enogene.com

Web: www.enogene.com