



NGFR Polyclonal Antibody

E92097

Catalog Number: E92097**Amount:** 100ul

Background: The p75 neurotrophin receptor (p75NTR), a member of the TNF receptor superfamily, is distinguished by multiple cysteine-rich ligand-binding domains, a single transmembrane sequence and a noncatalytic cytoplasmic domain (1). p75NTR displays paradoxical functions when acting alone or with other receptor proteins. Working in concert with Trk receptors, p75NTR recognizes neurotrophins and transmits trophic signals into the cell. Both p75NTR and TrkA are required to activate PI3K-Akt signaling, while TrkA can individually activate the MAP kinase pathway. In contrast, p75NTR, possibly through JNK, ensures appropriate apoptosis of injured neurons and improperly targeted neonatal neurons (2). The p75NTR protein undergoes sequential cleavage similar to APP and Notch. First, α -secretase removes the p75NTR ectodomain, eliminating ligand-mediated signaling. At this point, the membrane-tethered cleavage product can still fine-tune Trk-mediated trophic actions. γ -secretase cleaves within the transmembrane domain to liberate the cytoplasmic tail from its membrane anchor and allow the p75NTR intracellular domain to translocate to the nucleus (3,4).

Species: Rabbit**Isotype:** IgG

Storage/Stability: Store at -20°C or -80°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonyms: CD271; Gp80-LNGFR; TNFRSF16; p75(NTR); p75NTR;

Immunogen: A synthetic peptide of human NGFR

Purification: Affinity purification

Reactivity: H M R

Applications: WB IHC


Molecular Weight: 45kDa

Swiss-Prot No. : P08138

Gene ID: 4804

References: 1. Chao, M.V. (2003) Nat. Rev. Neurosci. 4, 299-309. 2. Nykjaer, A. et al. (2005) Curr. Opin. Neurobiol. 15, 49-57. 3. Kanning, K.C. et al. (2003) J. Neurosci. 23, 5425-5436. 4. Jung, K.M. et al. (2003) J. Biol. Chem. 278, 42161-42169.

For Research Use Only



Western blot analysis of extracts of various cell lines,
using NGFR antibody.