



Histone H3K79me1Polyclonal Antibody

E92367

- Catalog Number:** E92367
- Amount:** 100ul
- Background:** Modulation of chromatin structure plays an important role in the regulation of transcription in eukaryotes. The nucleosome, made up of DNA wound around eight core histone proteins (two each of H2A, H2B, H3, and H4), is the primary building block of chromatin (1). The amino-terminal tails of core histones undergo various post-translational modifications, including acetylation, phosphorylation, methylation, and ubiquitination (2-5). These modifications occur in response to various stimuli and have a direct effect on the accessibility of chromatin to transcription factors and, therefore, gene expression (6). In most species, histone H2B is primarily acetylated at Lys5, 12, 15, and 20 (4,7). Histone H3 is primarily acetylated at Lys9, 14, 18, 23, 27, and 56. Acetylation of H3 at Lys9 appears to have a dominant role in histone deposition and chromatin assembly in some organisms (2,3). Phosphorylation at Ser10, Ser28, and Thr11 of histone H3 is tightly correlated with chromosome condensation during both mitosis and meiosis (8-10). Phosphorylation at Thr3 of histone H3 is highly conserved among many species and is catalyzed by the kinase haspin. Immunostaining with phospho-specific antibodies in mammalian cells reveals mitotic phosphorylation at Thr3 of H3 in prophase and its dephosphorylation during anaphase (11).
- Species:** Rabbit
- Isotype:** IgG
- Storage/Stability:** Store at -20oC or -80oC. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
- Synonyms:** HIST1H3J; H3/j ;H3FJ ; Histone H3.1; Histone H3/a; Histone H3/b;Histone H3/c; Histone H3/d; Histone H3/f;Histone H3/h ; Histone H3/l;HistoneH3/j ;Histone H3/k; Histone H3/l;HIST3H3;
- Immunogen:** A synthetic methylated peptide corresponding to residues surrounding K79 of human histone H3
- Purification:** Affinity purification
- Reactivity:** H M ROther (Wide Range)
- Applications:** WBIHC IF IP ChIP
- Molecular Weight:** 15kDa
- Swiss-Prot No. :** Q16695
- Gene ID:** 8290
- References:** 1. Workman, J.L. and Kingston, R.E. (1998) Annu Rev Biochem 67, 545-79. 2. Hansen, J.C. et al. (1998) Biochemistry 37, 17637-41. 3. Strahl, B.D. and Allis, C.D. (2000) Nature 403, 41-5. 4. Cheung, P. et al. (2000) Cell 103, 263-71. 5. Bernstein, B.E. and Schreiber, S.L. (2002) Chem Biol 9, 1167-73. 6. Jaskelioff, M. and Peterson, C.L. (2003) Nat Cell Biol 5, 395-9. 7. Thorne, A.W. et al. (1990) Eur J Biochem 193, 701-13. 8. Hendzel, M.J. et al. (1997) Chromosoma 106, 348-60. 9. Goto, H. et al. (1999) J Biol Chem 274, 25543-9. 10. Preuss, U. et al. (2003) Nucleic Acids Res 31, 878-85. 11. Dai, J. et al. (2005) Genes Dev 19, 472-88.

For Research Use Only


WB 1:500 - 1:2000

IHC 1:50 - 1:200


IF 1:50 - 1:200

IP 1:50 - 1:200


ChIP 1:50 - 1:200



Western blot analysis of extracts of HeLa cellline and H3 protein expressed in E.coli., using H3K79me1 antibody.



Dot-blot analysis of all sorts of methylation peptides using H3K79me1 antibody.



Immunofluorescence analysis of 293T cell using H3K79me1 antibody.

For Research Use Only