



IHH Polyclonal Antibody

E92474

Catalog Number: E92474**Amount:** 100ul

Background: Hedgehog proteins (Hh) are secreted signaling proteins that play many roles during animal development. Aberrant Hh signaling activity can be associated with numerous birth defects and uncontrolled Hh pathway activation is linked to the development of several types of cancers (1-2). The three identified vertebrate Hh genes are Sonic (Shh), Indian (Ihh) and Desert (Dhh), all of which have distinct as well as overlapping roles (3-5). Hh proteins are synthesized as 45 kDa precursors that undergo auto-cleavage to generate a 19 kDa amino-terminal peptide (Hh-N) and a carboxy-terminal peptide (Hh-C). The amino-terminal peptide becomes covalently attached to a cholesterol molecule at its carboxy terminus and acetylated at its amino terminus. This doubly modified Hh-N peptide is released from cells and responsible for all known Hedgehog signaling activity (6).

Species: Rabbit**Isotype:** IgG

Storage/Stability: Store at -20oC or -80oC. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonyms: IHH;BDA1;HHG2;Indian hedgehog protein C-product;HHG-2;

Immunogen: Fusion protein of human IHH

Purification: Affinity purification

Reactivity: H M R

Applications: WB

Molecular Weight: 45kDa

Swiss-Prot No.: Q14623

Gene ID: 3549

References: 1. Ingham, P.W. and McMahon, A.P. (2001) Genes Dev. 15, 3059-3087. 2. McMahon, A.P. et al. (2003) Curr. Top Dev. Biol. 53, 1-114. 3. Zhang, X.M. et al. (2001) Cell 106, 781-792. 4. Adolphe, C. et al. (2004) Development 131, 5009-5019. 5. Pathi, S. et al. (2001) Mech. Dev. 106, 107-117. 6. Bijlsma, M.F. et al. (2004) Bioessays 26, 387-394.

For Research Use Only

