



## CD154/CD40L/TNFSF5

E21-I56

**Catalog Number:**E21-I56

**Amount:**10ug

**Alternative:**CD40 Ligand; CD40-L; T-Cell Antigen Gp39; TNF-Related Activation Protein; TRAP; Tumor Necrosis Factor Ligand Superfamily Member 5; CD154; CD40LG; CD40L; TNFSF5; TRAP

**Storage/Stability:**Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

**Background:**CD40 Ligand (CD40LG) is a type II transmembrane glycoprotein that belongs to the TNF superfamily. Like other TNF superfamily members, CD40LG exists as a trimer in membrane bound and soluble form, both of which are bioactive. CD40LG is a ligand for CD40; its ligation also initiates signal transduction in CD40LG expressing cells. CD40LG is a differentiation antigen that is expressed on the surface of T-cells. It stimulates B-cell proliferation and secretion of all immunoglobulin isotypes in the presence of cytokines. CD40LG has been shown to induce cytokine production and tumoricidal activity in peripheral blood monocytes. It also co-stimulates proliferation of activated T-cells and this is accompanied by the production of IFN-gamma, TNF-alpha, and IL2.

**Species:**Human

**Reconstitution:**Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin:**Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.

**Purity:**Greater than 95% as determined by reducing SDS-PAGE.

**Description:**Recombinant Human CD40 Ligand is produced by our Mammalian expression system and the target gene encoding Met113-Leu261 is expressed.


**Product State:**Lyophilized

**Ship Description:**The product is shipped at ambient temperature.

**Formulation:**Lyophilized from a 0.2 µm filtered solution of 20mM PB, 200mM NaCl, 0.1mM EDTA, pH 7.0.

**Expression system:**Human cells

**For Research Use Only**



**For Research Use Only**