




Fc γ RIIa/FCGR2A/CD32a

E21-S35

Catalog Number:E21-S35**Amount:**10ug**Alternate:**Low affinity immunoglobulin gamma Fc region receptor II-a; IgG Fc receptor II-a; CDw32; Fc-gamma RII-a; Fc-gamma-RIIa; FcRII-a; CD32; FCGR2A; FCG2; FCGR2A1; IGFR2**Storage/Stability:**Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.**Background:**Human Fc γ Rs are divided into three classes designated Fc γ RI (CD64), Fc γ RII (CD32), and Fc γ RIII (CD16), which generate multiple isoforms, are recognized. The activating-type receptor either has or associates non-covalently with an accessory subunit that has an immunoreceptor tyrosine-based activation motif (ITAM) in its cytoplasmic domain. Fc γ RI binds IgG with high affinity and functions during early immune responses, whereas Fc γ RII and RIII are low affinity receptors that recognize IgG as aggregates surrounding multivalent antigens during late immune responses. Human CD32, also known as Low affinity immunoglobulin γ Fc region receptor II-a (IgG Fc receptor II-a), Fc γ RII A or FCGR2A Protein, is expressed on cells of both myeloid and lymphoid lineages as well as on cells of non-hematopoietic origin. Associated with an ITAM-bearing adapter subunit, FcR γ , CD32a (Fc γ RII A) delivers an activating signal upon ligand binding, and results in the initiation of inflammatory responses including cytosis, phagocytosis, degranulation, and cytokine production. The responses can be modulated by signals from the co-expressed inhibitory receptors such as Fc γ RII B, and the strength of the signal is dependent on the ratio of expression of the activating and inhibitory receptors.**Species:**Human**Reconstitution:**Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.**Endotoxin:**Less than 0.1 ng/ μ g (1 IEU/ μ g) as determined by LAL test.**Purity:**Greater than 95% as determined by reducing SDS-PAGE.**Description:**Recombinant Human Low Affinity Immunoglobulin Gamma Fc Region Receptor II-A is produced by our Mammalian expression system and the target gene encoding Ala36-Ile218(His131Arg) is expressed with a 6His tag at the C-terminus.**Product State:**Lyophilized**Ship Description:**The product is shipped at ambient temperature.**Formulation:**Lyophilized from a 0.2 μ m filtered solution of PBS, pH7.4.**Expression system:**Human cells

For Research Use Only



For Research Use Only