## SMAD2 Mouse mAb

at +4°C short term. Store at -20°C long term. Avoid  thaw cycle.  d recombinant fragment of human SMAD2 expressed in  fi.  fluid  MADH2; MADR2; JV18-1; hMAD-2; hSMAD2; MGC22139;
e / thaw cycle.  ed recombinant fragment of human SMAD2 expressed in i.  e fluid
i. c fluid
n
MADH2; MADR2; JV18-1; hMAD-2; hSMAD2; MGC22139;
4440
rotein encoded by this gene belongs to the SMAD, a of proteins similar to the gene products of the phila gene 'mothers against decapentaplegic' (Mad) and elegans gene Sma. SMAD proteins are signal lucers and transcriptional modulators that mediate le signaling pathways. This protein mediates the signal transforming growth factor (TGF)-beta, and thus tes multiple cellular processes, such as cell proliferation, osis, and differentiation. This protein is recruited to the

## For Research Use Only

	beta receptors. The phosphorylation induces the dissociation of this protein with SARA and the association with the family member SMAD4. The association with SMAD4 is important for the translocation of this protein into the nucleus, where it binds to target promoters and forms a transcription repressor complex with other cofactors. This protein can also be phosphorylated by activin type 1 receptor kinase, and mediates the signal from the activin. Alternatively spliced transcript variants encoding the same protein have been observed.
Gene ID:	4087
Cellular localization:	Cytoplasm, Nucleus
Source:	Mouse
Antibody type:	Monclonal antibody
Isotype:	Mouse IgG1
Molecular Weight:	52kDa
Preservative:	Ascitic fluid containing 0.03% sodium azide.
Recommended Dilutions:	WB: 1/500 - 1/2000; IHC: 1/200 - 1/1000; ICC: 1/200 - 1/1000; FCM: 1/200 - 1/400; Elisa: 1/10000
Clone Number:	5G7