



Swiss-Prot No.:	P16234
Altename:	PDGFRA
Storage/Stability:	Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
Immunogen:	Synthetic peptide conjugated to KLH.
Purification:	Affinity purified
Reactivity:	Human,Rat,Mouse
Background:	<p>Tyrosine-protein kinase that acts as a cell-surface receptor for PDGFA, PDGFB and PDGFC and plays an essential role in the regulation of embryonic development, cell proliferation, survival and chemotaxis. Depending on the context, promotes or inhibits cell proliferation and cell migration. Plays an important role in the differentiation of bone marrow-derived mesenchymal stem cells. Required for normal skeleton development and cephalic closure during embryonic development. Required for normal development of the mucosa lining the gastrointestinal tract, and for recruitment of mesenchymal cells and normal development of intestinal villi. Plays a role in cell migration and chemotaxis in wound healing. Plays a role in platelet activation, secretion of agonists from platelet granules, and in thrombin-induced platelet aggregation. Binding of its cognate ligands - homodimeric PDGFA, homodimeric PDGFB, heterodimers formed by PDGFA and PDGFB or homodimeric PDGFC -leads</p>

to the activation of several signaling cascades; the response depends on the nature of the bound ligand and is modulated by the formation of heterodimers between PDGFRA and PDGFRB. Phosphorylates PIK3R1, PLCG1, and PTPN11. Activation of PLCG1 leads to the production of the cellular signaling molecules diacylglycerol and inositol 1,4,5-trisphosphate, mobilization of cytosolic Ca²⁺ and the activation of protein kinase C. Phosphorylates PIK3R1, the regulatory subunit of phosphatidylinositol 3-kinase, and thereby mediates activation of the AKT1 signaling pathway. Mediates activation of HRAS and of the MAP kinases MAPK1/ERK2 and/or MAPK3/ERK1. Promotes activation of STAT family members STAT1, STAT3 and STAT5A and/or STAT5B. Receptor signaling is down-regulated by protein phosphatases that dephosphorylate the receptor and its down-stream effectors, and by rapid internalization of the activated receptor.

Gene ID:	5156
Cellular localization:	Cell membrane, Membrane
Source:	Mouse
Antibody type:	Monclonal antibody
Isotype:	IgG1
Molecular Weight:	180kDa
Preservative:	PBS(pH 7.4) containing with 0.02% sodium azide and 50% glycerol.
Recommended Dilutions:	IHC 1:100-200 (Optimal dilutions should be determined by the end user)
Clone Number:	4G1-6A10-9C2

Form of Antibody:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
--------------------------	---