

	<h1>PIK3CA Rabbit pAb</h1>	E 2 5 0 0 2 6 5
--	----------------------------	--------------------------------------

<b>Swiss-Prot No.:</b>	P42336
<b>Altername:</b>	PIK3CA
<b>Storage/Stability:</b>	Store at -20°C. Avoid freeze / thaw cycles.
<b>Immunogen:</b>	Recombinant fusion protein containing a sequence corresponding to amino acids 401-600 of human PIK3CA (NP_006209.2).
<b>Purification:</b>	Affinity purified
<b>Reactivity:</b>	Human,Mouse,Rat
<b>Other Names:</b>	MCM; CWS5; MCAP; PI3K; CLOVE; MCMT C; PI3K-alpha; p110-alpha
	<p>Phosphoinositide-3-kinase (PI3K) that phosphorylates PtdIns (Phosphatidylinositol), PtdIns4P (Phosphatidylinositol 4-phosphate) and PtdIns(4,5)P2 (Phosphatidylinositol 4,5-bisphosphate) to generate phosphatidylinositol 3,4,5-trisphosphate (PIP3). PIP3 plays a key role by recruiting PH domain-containing proteins to the membrane, including AKT 1 and PDK1, activating signaling cascades involved in cell growth, survival, proliferation, motility and morphology. Participates in cellular signaling in response to various growth factors. Involved in the activation of AKT1 upon stimulation by receptor tyrosine kinases ligands such as EGF, insulin, IGF1, VEGFA and PDGF. Involved in signaling via insulin-receptor</p>

**For Research Use Only**

<b>Relevance:</b>	substrate (IRS) proteins. Essential in endothelial cell migration during vascular development through VEGFA signaling, possibly by regulating RhoA activity. Required for lymphatic vasculature development, possibly by binding to RAS and by activation by EGF and FGF2, but not by PDGF. Regulates invadopodia formation through the PDPK1-AKT1 pathway. Participates in cardiomyogenesis in embryonic stem cells through a AKT1 pathway. Participates in vasculogenesis in embryonic stem cells through PDK1 and protein kinase C pathway. Also has serine-protein kinase activity: phosphorylates PIK3R1 (p85alpha regulatory subunit), EIF4EBP1 and HRAS. Plays a role in the positive regulation of phagocytosis and pinocytosis .
<b>Source:</b>	Rabbit
<b>Antibody type:</b>	Polyclonal antibody
<b>Isotype:</b>	Rabbit IgG
<b>Molecular Weight:</b>	115-120kDa
<b>Preservative:</b>	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
<b>Recommended Dilutions:</b>	WB 1:500 - 1:2000; IHC 1:50 - 1:100(Optimal dilutions should be determined by the end user)