

	<h1>MSH6 Rabbit pAb</h1>	E 2 5 0 0 9 8 3
--	--------------------------	--------------------------------------

Swiss-Prot No.:	P52701
Altername:	MSH6
Storage/Stability:	Store at -20°C. Avoid freeze / thaw cycles.
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-290 of human MSH6 (NP_000170.1).
Purification:	Affinity purified
Reactivity:	Human,Mouse,Monkey
Other Names:	GTBP; HSAP; p160; GTMBP; HNPCC5
Cellular localization:	Chromosome, Nucleus
	Component of the post-replicative DNA mismatch repair system (MMR). Heterodimerizes with MSH2 to form MutS alpha, which binds to DNA mismatches thereby initiating DNA repair. When bound, MutS alpha bends the DNA helix and shields approximately 20 base pairs, and recognizes single base mismatches and dinucleotide insertion-deletion loops (IDL) in the DNA. After mismatch binding, forms a ternary complex with the MutL alpha heterodimer, which is thought to be responsible for directing the downstream MMR events, including strand discrimination, excision, and resynthesis. ATP binding and hydrolysis play a pivotal role in mismatch repair functions. The

For Research Use Only

Relevance:	ATPase activity associated with MutS alpha regulates binding similar to a molecular switch: mismatched DNA provokes ADP-->ATP exchange, resulting in a discernible conformational transition that converts MutS alpha into a sliding clamp capable of hydrolysis-independent diffusion along the DNA backbone. This transition is crucial for mismatch repair. MutS alpha may also play a role in DNA homologous recombination repair. Recruited on chromatin in G1 and early S phase via its PWWP domain that specifically binds trimethylated 'Lys-36' of histone H3 (H3K36me3): early recruitment to chromatin to be replicated allowing a quick identification of mismatch repair to initiate the DNA mismatch repair reaction.
Source:	Rabbit
Antibody type:	Polyclonal antibody
Isotype:	Rabbit IgG
Molecular Weight:	160kDa
Preservative:	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Recommended Dilutions:	WB 1:500 - 1:2000; IHC 1:50 - 1:200; IF 1:20 - 1:100 (Optimal dilutions should be determined by the end user)