



<b>Swiss-Prot No.:</b>	P04629
<b>Altename:</b>	NTRK1
<b>Storage/Stability:</b>	Store at -20°C. Avoid freeze / thaw cycles.
<b>Immunogen:</b>	Recombinant fusion protein containing a sequence corresponding to amino acids 1-280 of human NTRK1 (NP_002520.2).
<b>Purification:</b>	Affinity purified
<b>Reactivity:</b>	Human,Mouse,Rat
<b>Other Names:</b>	MTC; TRK; TRK1; TRKA; Trk-A; p140-TrkA
<b>Cellular localization:</b>	Cell membrane, Endosome, Membrane
	Receptor tyrosine kinase involved in the development and the maturation of the central and peripheral nervous systems through regulation of proliferation, differentiation and survival of sympathetic and nervous neurons. High affinity receptor for NGF which is its primary ligand (PubMed:1850821, PubMed:1849459, PubMed:1281417, PubMed:8325889, PubMed:15488758, PubMed:17196528). Can also bind and be activated by NTF3/neurotrophin-3. However, NTF3 only supports axonal extension through NTRK1 but has no effect on neuron survival . Upon dimeric NGF ligand-binding, undergoes homodimerization, autophosphorylation and

<b>Relevance:</b>	activation (PubMed:1281417). Recruits, phosphorylates and/or activates several downstream effectors including SHC1, FRS2, SH2B1, SH2B2 and PLCG1 that regulate distinct overlapping signaling cascades driving cell survival and differentiation. Through SHC1 and FRS2 activates a GRB2-Ras-MAPK cascade that regulates cell differentiation and survival. Through PLCG1 controls NF-Kappa-B activation and the transcription of genes involved in cell survival. Through SHC1 and SH2B1 controls a Ras-PI3 kinase-AKT1 signaling cascade that is also regulating survival. In absence of ligand and activation, may promote cell death, making the survival of neurons dependent on trophic factors.
<b>Source:</b>	Rabbit
<b>Antibody type:</b>	Polyclonal antibody
<b>Isotype:</b>	Rabbit IgG
<b>Molecular Weight:</b>	110kDa
<b>Preservative:</b>	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
<b>Recommended Dilutions:</b>	WB 1:500 - 1:2000; IHC 1:50 - 1:200; IF 1:50 - 1:200(Optimal dilutions should be determined by the end user)