

	<h1>TLR2 Rabbit pAb</h1>	E 2 5 0 2 5 4 5
--	--------------------------	--------------------------------------

<b>Swiss-Prot No.:</b>	O60603
<b>Altername:</b>	TLR2
<b>Storage/Stability:</b>	Store at -20°C. Avoid freeze / thaw cycles.
<b>Immunogen:</b>	Recombinant fusion protein containing a sequence corresponding to amino acids 19-300 of human TLR2 (NP_003255.2).
<b>Purification:</b>	Affinity purified
<b>Reactivity:</b>	Human
<b>Other Names:</b>	TIL4; CD282
<b>Cellular localization:</b>	Cytoplasmic vesicle, Membrane
	Cooperates with LY96 to mediate the innate immune response to bacterial lipoproteins and other microbial cell wall components. Cooperates with TLR1 or TLR6 to mediate the innate immune response to bacterial lipoproteins or lipopeptides (PubMed:21078852, PubMed:17889651). Acts via MYD88 and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response. May also activate immune cells and promote apoptosis in response to the lipid moiety of lipoproteins (PubMed:10426995, PubMed:10426996). Recognizes mycoplasmal macrophage-activating lipopeptide-2kD (MALP-2), soluble tuberculosis factor (STF), phenol-

**For Research Use Only**

<b>Relevance:</b>	soluble modulin (PSM) and B.burgdorferi outer surface protein A lipoprotein (OspA-L) cooperatively with TLR6 (PubMed:11441107). Stimulation of monocytes in vitro with M.tuberculosis PstS1 induces p38 MAPK and ERK1/2 activation primarily via this receptor, but also partially via TLR4 (PubMed:16622205). MAPK activation in response to bacterial peptidoglycan also occurs via this receptor (PubMed:16622205). Acts as a receptor for M.tuberculosis lipoproteins LprA, LprG, LpqH and PstS1, some lipoproteins are dependent on other coreceptors (TLR1, CD14 and/or CD36); the lipoproteins act as agonists to modulate antigen presenting cell functions in response to the pathogen (PubMed:19362712). M.tuberculosis HSP70 (dnaK) but not HSP65 (groEL-2) acts via this protein to stimulate NF-kappa-B expression (PubMed:15809303). Recognizes M.tuberculosis major T-antigen EsxA (ESAT-6) which inhibits downstream MYD88-dependent signaling (shown in mouse) . Forms activation clusters composed of several receptors depending on the ligand, these clusters trigger signaling from the cell surface and subsequently are targeted to the Golgi in a lipid-raft dependent pathway. Forms the cluster TLR2:TLR6:CD14:CD36 in response to diacylated lipopeptides and TLR2:TLR1:CD14 in response to triacylated lipopeptides (PubMed:16880211). Required for normal uptake of M.tuberculosis, a process that is inhibited by M.tuberculosis LppM .
<b>Source:</b>	Rabbit
<b>Antibody type:</b>	Polyclonal antibody
<b>Isotype:</b>	Rabbit IgG
<b>Molecular Weight:</b>	100kDa
<b>Preservative:</b>	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**For Research Use Only**

Order: [order@enogene.com](mailto:order@enogene.com) Support: [service@enogene.com](mailto:service@enogene.com) Web:  
[www.enogene.com](http://www.enogene.com)

---

<b>Recommended Dilutions:</b>	WB 1:500 - 1:2000 (Optimal dilutions should be determined by the end user)
-------------------------------	--

---

**For Research Use Only**