

NDE1 Rabbit pAb

| Swiss-Prot No.: | Q9NXR1 |
|---------------------------|---|
| Altername: | NDE1 |
| Storage/Stability: | Store at -20°C. Avoid freeze / thaw cycles. |
| Immunogen: | Recombinant fusion protein containing a sequence corresponding to amino acids 1-110 of human NDE1 (NP_060138.1). |
| Purification: | Affinity purified |
| Reactivity: | Human,Mouse,Rat |
| Other Names: | NDE; LIS4; MHAC; NUDE; NUDE1; HOM-TES-87 |
| Cellular localization: | Centromere, Chromosome, Cytoplasm, Cytoskeleton, Kinetochore, Microtubule |
| Relevance: | Required for centrosome duplication and formation and function of the mitotic spindle. Essential for the development of the cerebral cortex. May regulate the production of neurons by controlling the orientation of the mitotic spindle during division of cortical neuronal progenitors of the proliferative ventricular zone of the brain. Orientation of the division plane perpendicular to the layers of the cortex gives rise to two proliferative neuronal progenitors whereas parallel orientation of the division plane yields one proliferative neuronal progenitor and a post-mitotic neuron. A premature shift towards a neuronal fate within the progenitor population may |

For Research Use Only

Order: order@enogene.com Support: service@enogene.com Web: www.enogene.com

| | result in an overall reduction in the final number of neurons and an increase in the number of neurons in the deeper layers of the cortex. |
|---------------------------|--|
| Source: | Rabbit |
| Antibody type: | Polyclonal antibody |
| Isotype: | Rabbit IgG |
| Molecular Weight: | 39kDa |
| Preservative: | PBS with 0.02% sodium azide, 50% glycerol, pH7.3. |
| Recommended Dilutions: | WB 1:500 - 1:2000; IHC 1:50 - 1:200(Optimal dilutions should be determined by the end user) |