

	<h1 style="text-align: center;">XIAP Rabbit pAb</h1>	<div>E 2 5 1 3 2 7 7</div>
--	--	--

Swiss-Prot No.:	P98170
Altername:	XIAP
Storage/Stability:	Store at -20°C. Avoid freeze / thaw cycles.
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-200 of human XIAP (NP_001158.2).
Purification:	Affinity purified
Reactivity:	Human,Mouse
Other Names:	API3; ILP1; MIHA; XLP2; BIRC4; IAP-3; hIAP3; hIAP-3
Cellular localization:	Cytoplasm, Nucleus
	<p>Multi-functional protein which regulates not only caspases and apoptosis, but also modulates inflammatory signaling and immunity, copper homeostasis, mitogenic kinase signaling, cell proliferation, as well as cell invasion and metastasis. Acts as a direct caspase inhibitor. Directly bind to the active site pocket of CASP3 and CASP7 and obstructs substrate entry.</p> <p>Inactivates CASP9 by keeping it in a monomeric, inactive state.</p> <p>Acts as an E3 ubiquitin-protein ligase regulating NF-kappa-B signaling and the target proteins for its E3 ubiquitin-protein ligase activity include: RIPK1, CASP3, CASP7, CASP8, CASP9, MAP3K2/MEKK2, DIABLO/SMAC, AIFM1, CCS and</p>

For Research Use Only

Relevance:	<p>BIRC5/survivin. Ubiquitination of CCS leads to enhancement of its chaperone activity toward its physiologic target, SOD1, rather than proteasomal degradation. Ubiquitination of MAP3K2/MEKK2 and AIFM1 does not lead to proteasomal degradation. Plays a role in copper homeostasis by ubiquitinating COMMD1 and promoting its proteasomal degradation. Can also function as E3 ubiquitin-protein ligase of the NEDD8 conjugation pathway, targeting effector caspases for neddylation and inactivation. Regulates the BMP signaling pathway and the SMAD and MAP3K7/TAK1 dependent pathways leading to NF-kappa-B and JNK activation. Acts as an important regulator of innate immune signaling via regulation of Nodlike receptors (NLRs). Protects cells from spontaneous formation of the ripoptosome, a large multi-protein complex that has the capability to kill cancer cells in a caspase-dependent and caspase-independent manner. Suppresses ripoptosome formation by ubiquitinating RIPK1 and CASP8. Acts as a positive regulator of Wnt signaling and ubiquitinates TLE1, TLE2, TLE3, TLE4 and AES. Ubiquitination of TLE3 results in inhibition of its interaction with TCF7L2/TCF4 thereby allowing efficient recruitment and binding of the transcriptional coactivator beta-catenin to TCF7L2/TCF4 that is required to initiate a Wnt-specific transcriptional program.</p>
Source:	Rabbit
Antibody type:	Polyclonal antibody
Isotype:	Rabbit IgG
Molecular Weight:	51kDa
Preservative:	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Recommended	WB 1:500 - 1:2000; IF 1:50 - 1:200 (Optimal dilutions should be

For Research Use Only

Dilutions:	determined by the end user)
-------------------	-----------------------------