



|                               |                                                                                                                                           |
|-------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Altername:</b>             | ; NR1B1; RAR;                                                                                                                             |
| <b>Immunogen:</b>             | A peptide corresponding to the C terminal residues of human RARA protein.                                                                 |
| <b>Purification:</b>          | Purified by protein A/G column.                                                                                                           |
| <b>Reactivity:</b>            | Human                                                                                                                                     |
| <b>References:</b>            | Zhu HH, Qin YZ and Huang XJ. Resistance to arsenic therapy in acute promyelocytic leukemia. <i>N Engl J Med.</i> 370(19):1864-1866 (2014) |
| <b>Background:</b>            | Retinoic acid receptor alpha                                                                                                              |
| <b>Gene ID:</b>               | P10276                                                                                                                                    |
| <b>Cellular localization:</b> | Cytoplasm, Nucleus                                                                                                                        |
| <b>Source:</b>                | Rabbit                                                                                                                                    |
| <b>Antibody type:</b>         | Polyclonal                                                                                                                                |
| <b>Isotype:</b>               | IgG                                                                                                                                       |
| <b>Molecular Weight:</b>      | 51KD                                                                                                                                      |
| <b>Preservative:</b>          | Already dissolved in sterile H <sub>2</sub> O with 1% BSA and 0.02% NaN <sub>3</sub> and 50% glycerol.                                    |
| <b>Recommended</b>            | ELISA(1:20,000~80,000) Western blotting(1:200~1,000)                                                                                      |

**Dilutions:**