



<b>Altername:</b>	; PKCE; nPKC-epsilon;
<b>Immunogen:</b>	Recombinant fragment corresponding to 353-512aa of human PRKCE.
<b>Purification:</b>	Purified by antigen affinity column.
<b>Reactivity:</b>	Human. Predicted: Rat, Mus.
<b>References:</b>	Sun Z, Guo YS, Yan SJ, et al. CK8 phosphorylation induced by compressive loads underlies the downregulation of CK8 in human disc degeneration by activating protein kinase C. <i>Lab Invest.</i> 93 (12):1323-1330 (2013)
<b>Background:</b>	Homo sapiens protein kinase C, epsilon (PRKCE), mRNA.
<b>Gene ID:</b>	BC109033
<b>Cellular localization:</b>	Cytoplasm, Membrane, Nucleus
<b>Source:</b>	Rabbit
<b>Antibody type:</b>	Polyclonal
<b>Isotype:</b>	IgG
<b>Molecular Weight:</b>	84KD
<b>Preservative:</b>	Already dissolved in sterile H <sup>2</sup> O with 1% BSA and 0.02% NaN <sup>3</sup> and 50% glycerol.

<b>Recommended Dilutions:</b>	ELISA(1:20,000~80,000) Western blotting(1:200~1,000) Immunohistochemistry(1:100~500)
-------------------------------	---