

	<p>SLU7</p>	<p>E 3 0 8 1 9 4</p>
--	-------------	--

<b>Altername:</b>	; 9G8; hSlu7;
<b>Immunogen:</b>	Recombinant fragment corresponding to 297-488aa of human SLU7.
<b>Purification:</b>	Purified by antigen affinity column.
<b>Reactivity:</b>	Human. Predicted: Rat, Mus.
<b>References:</b>	Banerjee S, Khandelia P, Melangath G, et al. Splicing functions and global dependency on fission yeast slu7 reveal diversity in spliceosome assembly. Mol Cell Biol. 33(16):3125-3136 (2013)
<b>Background:</b>	Pre-mRNA-splicing factor SLU7
<b>Gene ID:</b>	BC010634
<b>Cellular localization:</b>	Cytoplasm, Nucleus, Spliceosome
<b>Source:</b>	Rabbit
<b>Antibody type:</b>	Polyclonal
<b>Isotype:</b>	IgG
<b>Molecular Weight:</b>	68KD
<b>Preservative:</b>	Already dissolved in sterile H <sub>2</sub> O with 1% BSA and 0.02% NaN <sub>3</sub> and 50% glycerol.
<b>Recommended</b>	ELISA(1:20,000~80,000) Western blotting(1:200~1,000)

**For Research Use Only**

Order: [order@enogene.com](mailto:order@enogene.com) Support: [service@enogene.com](mailto:service@enogene.com) Web:  
[www.enogene.com](http://www.enogene.com)

---

<b>Dilutions:</b>	
-------------------	--

---

**For Research Use Only**