

	<h1>PCBP4</h1>	E 3 1 0 0 4 0
--	----------------	---------------------------------

<b>Altername:</b>	; CBP; LIP4; MCG10;
<b>Immunogen:</b>	Recombinant fragment corresponding to 9-173aa of human PCBP4.
<b>Purification:</b>	Purified by antigen affinity column.
<b>Reactivity:</b>	Human. Predicted: Rat, Mus.
<b>References:</b>	Zhu J, Chen X. MCG10, a novel p53 target gene that encodes a KH domain RNA-binding protein, is capable of inducing apoptosis and cell cycle arrest in G2-M[J]. Molecular and Cellular Biology. 20(15): 5602-5618 (2000)
<b>Background:</b>	Homo sapiens poly(rC) binding protein 4(PCBP4), mRNA.
<b>Gene ID:</b>	BC017098
<b>Cellular localization:</b>	Cytoplasm
<b>Source:</b>	Rabbit
<b>Antibody type:</b>	Polyclonal
<b>Isotype:</b>	IgG
<b>Molecular Weight:</b>	44KD
<b>Preservative:</b>	Already dissolved in sterile H <sup>2</sup> O with 1% BSA and 0.02% NaN <sup>3</sup> and 50% glycerol.

**For Research Use Only**

Order: [order@enogene.com](mailto:order@enogene.com) Support: [service@enogene.com](mailto:service@enogene.com) Web:  
[www.enogene.com](http://www.enogene.com)

---

<b>Recommended Dilutions:</b>	ELISA (1:20,000-80,000) Western blotting (1:200-1,000)
-----------------------------------	--

---

**For Research Use Only**