


| | | |
|--|--------------|--|
|  | PDRG1 | E 3 1 0 0 5 4 |
|--|--------------|--|

| | |
|-------------------------------|--|
| Altername: | ; PDRG; |
| Immunogen: | Recombinant fragment corresponding to 12-126aa of human PDRG1. |
| Purification: | Purified by antigen affinity column. |
| Reactivity: | Human. Predicted: Rat, Mus. |
| References: | Bonaldo MF, Lennon G, Soares MB. Normalization and subtraction: two approaches to facilitate gene discovery. Genome Res. 6(9):791-806 (1996) |
| Background: | Homo sapiens p53 and DNA damage regulated 1, mRNA. |
| Gene ID: | BC009334 |
| Cellular localization: | Cytoplasm |
| Source: | Rabbit |
| Antibody type: | Polyclonal |
| Isotype: | IgG |
| Molecular Weight: | 16KD |
| Preservative: | Already dissolved in sterile H ₂ O with 1% BSA and 0.02% NaN ₃ and 50% glycerol. |
| Recommended | ELISA (1:20,000-80,000) Western blotting (1:500-1,000) |

For Research Use Only

Order: order@enogene.com Support: service@enogene.com Web:
www.enogene.com

| | |
|-------------------|----------------------------------|
| Dilutions: | Immunohistochemistry (1:100-500) |
|-------------------|----------------------------------|

For Research Use Only