SQSTM1

Altername:	; A170; DMRV; FTDALS3; NADGP; OSIL; PDB3; ZIP3; p60; p62; p62B;
Immunogen:	Recombinant fragment corresponding to 257-440aa of human SQSTM1.
Purification:	Purified by antigen affinity column.
Reactivity:	Human. Predicted: Rat, Mus.
References:	Park I, Chung J, Walsh CT, et al. Phosphotyrosine-independent binding of a 62-kDa protein to the src homology 2 (SH2) domain of p56lck and its regulation by phosphorylation of Ser-59 in the lck unique N-terminal region. Proc. Natl. Acad. Sci. U.S.A. 92 (26640,16639,16638,16637)
Other Names:	Purified Rabbit Anti Human SQSTM1 Polyclonal antibody.
Background:	Homo sapiens sequestosome 1, mRNA
Gene ID:	BC003139
Cellular localization:	Cytoplasm,?Cytoplasmic vesicle,?Endoplasmic reticulum,? Endosome,?Lysosome,?Nucleus
Source:	Rabbit
Antibody type:	Polyclonal
Isotype:	lgG
Molecular	62KD

For Research Use Only

Order: order@enogene.com Support: service@enogene.com Web: www.enogene.com

Weight:	
Preservative:	Already dissolved in sterile H^2O with 1% BSA and 0.02% NaN^3 and 50% glycerol.
Recommended Dilutions:	ELISA (1:20,000-80,000) Western blotting (1:200-1,000) Immunohistochemistry (1:100-500)