

	<h1>VEGFA</h1>	E 3 1 2 2 4 1
----------------------------------------------------------------------------------	----------------	----------------------------------------------------------------------------------

Altername:	; MVCD1; VEGF; VPF;
Immunogen:	Recombinant fragment corresponding to 177-368aa of human VEGFA.
Purification:	Purified by protein A/G column.
Reactivity:	Human. Predicted: Rat, Mus.
References:	Guan H, Zhou Z, Cao Y. et al. VEGF165 promotes the osteolytic bone destruction of ewing's sarcoma tumors by upregulating RANKL. Oncol Res. 18(2-3):117-125 (2009)
Background:	Homo sapiens vascular endothelial growth factor A (VEGFA)
Gene ID:	NM_001025368
Cellular localization:	Cytoplasmic and Secreted
Source:	Rabbit
Antibody type:	Polyclonal
Isotype:	IgG
Molecular Weight:	47KD
Preservative:	Already dissolved in sterile H ₂ O with 1% BSA and 0.02% NaN ₃ and 50% glycerol.
Recommended	ELISA(1:10,000~80,000) Western blotting (1:200~1,000)

For Research Use Only

Order: order@enogene.com Support: service@enogene.com Web:
www.enogene.com

Dilutions:	Immunohistochemistry (1:50~200) Immunofluoresc
-------------------	------------------------------------------------

For Research Use Only