


| | | |
|--|---|--|
|  | <p style="text-align: center;">YBX1</p> | <p style="text-align: center;">E 3 1 2 9 2 2</p> |
|--|---|--|

| | |
|-------------------------------|---|
| Altername: | ; YB1; BP-8; CSDB; DBPB; YB-1; CBF-A; CSDA2; EFI-A; NSEP1; NSEP-1; MDR-NF1; |
| Immunogen: | Recombinant fragment corresponding to 281-299aa of human YBX1. |
| Purification: | Purified by antigen affinity column. |
| Reactivity: | Human. Predicted:Rat, Mus. |
| References: | Shurtleff M J, Yao J, Qin Y. et al. Broad role for YBX1 in defining the small noncoding RNA composition of exosomes. Proceedings of the National Academy of Sciences. 114 (43):E8987-E8995 (2017) |
| Other Names: | Purified Rabbit Anti Human YBX1 Polyclonal antibody. |
| Background: | Homo sapiens Y-box binding protein 1 (YBX1), transcript variant 1, mRNA |
| Gene ID: | NM_004559 |
| Cellular localization: | Cytoplasm, Nucleus, Secreted |
| Source: | Rabbit |
| Antibody type: | Polyclonal |
| Isotype: | IgG |
| Molecular | 50KD |

For Research Use Only

| | |
|-------------------------------|--|
| Weight: | |
| Preservative: | Already dissolved in sterile H ₂ O with 1% BSA and 0.02% NaN ³ and 50% glycerol. |
| Recommended Dilutions: | ELISA (1:20,000-80,000) Western blotting (1:200-1,000) |