



Antibody type:	Polyclonal Antibody
Applications:	WB
Reactivity:	Human Mouse Rat
Molecular Weight:	48kDa
Immunogen:	A synthetic peptide of human SKP2
Gene ID:	6502
Swiss-Prot No.:	Q13309
Alternname:	FBL1;FBXL1;FLB1;p45
Source:	Rabbit
Isotype:	IgG
Purification:	Affinity purification
Storage/Stability:	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
	This gene encodes a member of the F-box protein family which is characterized by an approximately 40 amino acid motif, the F-box. The F-box proteins constitute one of the four subunits of ubiquitin protein ligase complex called SCFs (SKP1-cullin-F-box), which function in phosphorylation-dependent ubiquitination. The F-box proteins are divided into 3 classes: Fbws containing WD-40 domains, Fbls containing leucine-rich repeats, and Fbxs containing either different

Background:	protein-protein interaction modules or no recognizable motifs. The protein encoded by this gene belongs to the Fbls class; in addition to an F-box, this protein contains 10 tandem leucine-rich repeats. This protein is an essential element of the cyclin A-CDK2 S-phase kinase. It specifically recognizes phosphorylated cyclin-dependent kinase inhibitor 1B (CDKN1B, also referred to as p27 or KIP1) predominantly in S phase and interacts with S-phase kinase-associated protein 1 (SKP1 or p19). In addition, this gene is established as a protooncogene causally involved in the pathogenesis of lymphomas. Alternative splicing of this gene generates three transcript variants encoding different isoforms.
Dilution:	WB 1:500 - 1:2000
Shipping&Stablity:	Aliquot and store at -20°C. Avoid repeated freeze / thaw cycles.